

Bulk Boundary-Layer Models

David A. Randall

*Department of Atmospheric Science
Colorado State University, Fort Collins, Colorado 80523*

Ball (1960) was the first to propose a model in which the interior of the planetary boundary layer (PBL) is well-mixed in the conservative variables, while the PBL top is marked by discontinuities in these same variables. Geisler and Kraus (1969) were the first to extend the idea by treating momentum (not a conservative variable) in the same framework, so that the PBL could be said to move as a “slab.” There is now a huge literature on mixed layer models, and in almost every paper the emphasis is on the question of what determines the time rate of change of the PBL depth. Actually, many of the papers are devoted to the *ocean* mixed layer, which is a kind of upside-down PBL. More generally, any model in which the PBL depth is an explicit parameter and the vertical structure of the PBL is described with just a few degrees of freedom can be called a “bulk” PBL model.

We begin our study of bulk models by deriving the equations that govern the vertically-averaged properties of the PBL. Let A be an arbitrary intensive scalar, satisfying the “flux-form” conservation equation

$$\frac{\partial}{\partial t} (\rho A) + \nabla \cdot (\rho \mathbf{V} A) + \frac{\partial}{\partial z} (\rho w A) = - \frac{\partial F_A}{\partial z} + S_A , \quad (1)$$

where $F_A \equiv \overline{\rho w' A'}$ is the upward turbulent flux of A , bars are omitted on the mean quantities, and S_A is a source or sink of A , per unit volume. The corresponding continuity equation is

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{V}) + \frac{\partial}{\partial z} (\rho w) = 0 , \quad (2)$$

which can also be written as

$$\frac{\partial}{\partial t} \rho + \mathbf{V} \cdot \nabla \rho + w \frac{\partial}{\partial z} \rho = -\rho \left(\nabla \cdot \mathbf{V} + \frac{\partial}{\partial z} w \right) . \quad (3)$$

By use of (2), we can rewrite (1) in the “advective form:”

$$\rho \left(\frac{\partial A}{\partial t} + \mathbf{V} \cdot \nabla A + w \frac{\partial A}{\partial z} \right) = - \frac{\partial F_A}{\partial z} + S_A . \quad (4)$$

Integrating (1) from just below to just above the PBL top, and using Leibniz' rule, we get

$$\begin{aligned} \frac{\partial}{\partial t} \int_{z_B - \epsilon}^{z_B + \epsilon} \rho A dz - \Delta(\rho A) \frac{\partial z_B}{\partial t} + \nabla \cdot \int_{z_B - \epsilon}^{z_B + \epsilon} \rho \mathbf{V} A dz - \Delta(\rho \mathbf{V} A) \cdot \Delta z_B \\ + \Delta(\rho w A) = -(F_A)_{B+} + \int_{z_B - \epsilon}^{z_B + \epsilon} S_A dz , \end{aligned} \quad (5)$$

where the red terms drop out as the domain of integration shrinks to zero and/or because all of the turbulence variables go to zero above the PBL top. Here we have used the notation $\Delta(\cdot) \equiv (\cdot)_{z=z_B + \epsilon} - (\cdot)_{z=z_B - \epsilon} \equiv (\cdot)_{B+} - (\cdot)_B$, and henceforth subscripts $B+$ and B denote levels just above and just below the PBL top, respectively. For $A \equiv 1$, (5) reduces to mass conservation in the form

$$\rho_{B+} \left(\frac{\partial z_B}{\partial t} + \mathbf{V}_{B+} \cdot \nabla z_B - w_{B+} \right) = \rho_B \left(\frac{\partial z_B}{\partial t} + \mathbf{V}_B \cdot \nabla z_B - w_B \right) = E - M_B , \quad (6)$$

where $E - M_B$ is the total mass flux across the PBL top. In essence, (6) simply says that the mass flux is continuous across the PBL top, i.e., no mass is created or destroyed between levels B and $B+$. We interpret M_B as the mass flux due to a loss of PBL mass into cumulus clouds, and E as the mass flux due to the turbulent *entrainment* of free atmospheric air into the PBL. See Fig. 1 for a sketch illustrating the physical system under consideration. With the use of (6), we can rewrite (5) as

$$-\Delta A (E - M_B) = (F_A)_B + \int_{z_B - \epsilon}^{z_B + \epsilon} S_A dz , \quad (7)$$

Here we have assumed that $(F_A)_{B+} = 0$. For $S_A \equiv 0$, (7) simply says that the total flux of A must be continuous across the PBL top. Notice that for $\Delta A \neq 0$, a mass flux across the PBL top is generally associated with a turbulent flux of A at level B . This flux serves to change the A of entering particles from A_{B+} to A_B . For example, dry entrained air is moistened by an upward moisture flux which converges "discontinuously" at level B . Lilly (1968) was the first to derive (7) using the approach followed here.




Figure 1: Sketch of a column of PBL air, with arrows indicating the lateral fluxes due to the horizontal winds, and the top-flux due to entrainment and the cumulus mass flux.

Now integrate (1) through the PBL depth, from the surface to level B , to obtain

$$\begin{aligned} & \frac{\partial}{\partial t} \int_{z_S}^{z_B} (\rho A) dz - (\rho A)_B \frac{\partial z_B}{\partial t} + \\ & (\rho A)_S \frac{\partial z_S}{\partial t} + \nabla \cdot \int_{z_S}^{z_B} (\rho \mathbf{V} A) dz - (\rho \mathbf{V} A)_S \cdot \nabla z_S \\ & - (\rho \mathbf{V} A)_B \cdot \nabla z_B + (\rho w A)_B - (\rho w A)_S = (F_A)_S - (F_A)_B + \int_{z_S}^{z_B - \varepsilon} S_A dz . \end{aligned} \quad (8)$$

The condition that no mass crosses the Earth's surface can be written as

$$\rho_S \left(\frac{\partial z_S}{\partial t} + \mathbf{V}_S \cdot \nabla_{z_S} w_S \right) = 0 . \quad (9)$$

Use of (6) and (9) allows us to simplify (8) considerably, to

$$\begin{aligned} \frac{\partial}{\partial t} (\rho_M A_M \delta z_M) + \nabla \cdot [\rho_M (\mathbf{V} A)_M \delta z_M] - A_B (E - M_B) \\ = (F_A)_S - (F_A)_M + (S_A)_M \delta z_M . \end{aligned} \quad (10)$$

Here we define the depth of the PBL as

$$\delta z_M \equiv z_B - z_S \quad (11)$$

and

$$(S_A)_M \delta z_m \equiv \int_{z_S}^{z_B - \varepsilon} S_A dz . \quad (12)$$

By combining (7) with (10), we obtain

$$\begin{aligned} \frac{\partial}{\partial t} (\rho_M A_M \delta z_M) + \nabla \cdot [\rho_M (\mathbf{V} A)_M \delta z_M] - A_{B+} (E - M_B) \\ = (F_A)_S - (S_A)_M \delta z_m + \int_{z_B - \varepsilon}^{z_B + \varepsilon} S_A dz . \end{aligned} \quad (13)$$


For $A \equiv 1$, both (10) and (13) reduce to a statement of mass conservation for the whole PBL:

$$\frac{\partial}{\partial t} (\rho_M \delta z_M) + \nabla \cdot [(\rho \mathbf{V})_M \delta z_M] = E - M_B . \quad (14)$$

Now define a transformed vertical coordinate that follows the PBL top, given by

$$z' \equiv z - z_B(x, y, t) . \quad (15)$$

Fig. 2 illustrates the transformation of derivatives that is associated with this coordinate, as used below. We can write



$$\frac{A_2 - A_1}{\delta x} = \frac{A_3 - A_1}{\delta x} - \left(\frac{A_3 - A_2}{\delta z} \right) \frac{\delta z}{\delta x}$$

$$\left(\frac{\partial A}{\partial x} \right)_z = \left(\frac{\partial A}{\partial x} \right)_{z'} - \frac{\partial A}{\partial z} \frac{\partial z_B}{\partial x}$$

Figure 2: Sketch used in analyzing the coordinate transformation from constant z to constant z' .

$$\left(\frac{\partial \rho}{\partial t}\right)_z = \left(\frac{\partial \rho}{\partial t}\right)_{z'} - \frac{\partial \rho}{\partial z} \frac{\partial z_B}{\partial t}, \quad (16)$$

$$\nabla \cdot (\rho \mathbf{V}) = \nabla_{z'} \cdot (\rho \mathbf{V}) - \frac{\partial}{\partial z} (\rho \mathbf{V}) \nabla z_B. \quad (17)$$

Substitution of (16) and (17) into (2) gives

$$\left(\frac{\partial \rho}{\partial t}\right)_{z'} + \nabla_{z'} \cdot (\rho \mathbf{V}) + \frac{\partial(\rho w')}{\partial z} = 0, \quad (18)$$

where we define

$$w'(z) \equiv -\left[\frac{\partial z_B}{\partial t} + \mathbf{V}(z) \cdot \nabla z_B - w(z)\right]. \quad (19)$$

Note that w' is a function of height, and that

$$\rho_B w'(z_B) = \rho_{B+} w'(z_{B+}) = -(E - M_B). \quad (20)$$

Similarly, Eq. (1) can be rewritten as

$$\left[\frac{\partial(\rho A)}{\partial t}\right]_{z'} + \nabla_{z'} \cdot (\rho \mathbf{V} A) \frac{\partial(\rho w' A)}{\partial z'} = -\frac{\partial F_A}{\partial z} + S_A. \quad (21)$$

By combining (18) and (21), we obtain

$$\left(\frac{\partial A}{\partial t}\right)_{z'} + \mathbf{V} \cdot \nabla_{z'} A + w' \frac{\partial A}{\partial z} = -\alpha \frac{\partial F_A}{\partial z'} + \alpha S_A. \quad (22)$$

Equation (22) describes the time-rate-of-change of A as seen on any surface of constant z' . One such surface is the top of the boundary layer, so we can apply (22) to determine the time rate of change of a quantity on the surface $z = z_B(x, y, t)$. Evaluating (22) at levels $B+$ and B , subtracting, and using (6), we find that

$$\frac{\partial(\Delta A)}{\partial t} + \Delta(\mathbf{V} \cdot \nabla_{z'} A) - (E - M_B) \Delta\left(\alpha \frac{\partial A}{\partial z}\right) = a_B \left(\frac{\partial F_A}{\partial z}\right)_B + \Delta(\alpha S_A). \quad (23)$$

This governs the time rates of change of the ‘‘jumps.’’ In writing (23), we have assumed that the turbulent flux divergence vanishes at level $B+$.

Consider the special case of the horizontal momentum equation, i.e., $A \rightarrow \mathbf{V}$. Then we have $\alpha S_A \rightarrow -\alpha \nabla_z p - f \mathbf{k} \times \mathbf{V}$, where p is pressure and f is the Coriolis parameter. The gradient operator satisfies

$$\begin{aligned}
-\alpha \nabla_z p &= -\alpha \left(\nabla_{z'} p - \frac{\partial p}{\partial z} \nabla z_B \right) \\
&= \alpha (\nabla_{z'} p + \rho g \nabla z_B) \\
&= -\alpha \nabla_{z'} p - g \nabla z_B .
\end{aligned} \tag{24}$$

Here we have used hydrostatics. Applying (23) at levels $B+$ and B , and subtracting, we find that

$$-\Delta(\alpha \nabla_z p) = -\Delta\alpha \nabla p_B . \tag{25}$$

Then, corresponding to (23), we obtain

$$\begin{aligned}
\frac{\partial(\Delta \mathbf{V})}{\partial t} + \Delta(\mathbf{V} \cdot \nabla_{z'} A) - (E - M_B) \nabla \left(\alpha \frac{\partial \mathbf{V}}{\partial z} \right) = \\
\alpha_B \left(\frac{\partial \mathbf{F}_v}{\partial z} \right)_B - \Delta\alpha \nabla p_B - f \mathbf{k} \times \Delta \mathbf{V} .
\end{aligned} \tag{26}$$

For a steady state in which advection and friction are negligible, this reduces to

$$0 = -\Delta\alpha \nabla p_B - f \mathbf{k} \times \Delta \mathbf{V} , \tag{27}$$

which we recognize as a form of the thermal wind equation, sometimes called Margules' equation.

References and Bibliography

- Ball, F. K., 1960: Control of inversion height by surface heating. *Quart J. Roy. Meteor. Soc.*, **86**, 483–494.
- Geisler, J. E., & Kraus, E. B., 1969: The well-mixed Ekman boundary layer. *Deep Sea Res.*, **16** (Suppl. to Vo.), 73–84.
- Kraus, E. B., and J. S. Turner, 1967: A one-dimensional model of the seasonal thermocline. II. The general theory and its consequences. *Tellus*, **19**, 98-105.
- Lilly, D. K., 1968: Models of cloud-topped mixed layers under a strong inversion. *Quart. J. Roy. Meteor. Soc.*, **94**, 292–309.
- Randall, D. A., 1984: Buoyant production and consumption of turbulence kinetic energy in cloud-topped mixed layers. *J. Atmos. Sci.*, **41**, 402-413.
- Randall, D. A., and M. J. Suarez, 1984: On the dynamics of stratocumulus formation and dissipation. *J. Atmos. Sci.*, **41**, 3052-3057.
- Randall, D. A., 1985: Practical considerations concerning the use of the planetary boundary layer top as a coordinate surface. Technical Memo 103, European Centre for Medium Range Weather Forecasts, **12** pp.
- Randall, D. A., J. A. Abeles, and T. G. Corsetti, 1985: Seasonal simulations of the planetary boundary layer and boundary-layer stratocumulus clouds with a general circulation model. *J. Atmos. Sci.*, **42**, 641-676.
- Schubert, W. H., 1976: Experiments with Lilly's cloud-topped mixed layer model. *J. Atmos. Sci.*, **33**, 436-446.
- Schubert, W. H., J. S. Wakefield, E. J. Steiner, and S. K. Cox, 1979: Marine stratocumulus convection, Part I: Governing equations and horizontally homogeneous solutions. *J. Atmos. Sci.*, **36**, 1286-1307.
- Schubert, W. H., J. S. Wakefield, E. J. Steiner, and S. K. Cox, 1979: Marine stratocumulus convection. Part II: Horizontally inhomogeneous solutions. *J. Atmos. Sci.*, **36**, 1308-1324.

Suarez, M. J., A. Arakawa, and D. A. Randall, 1983: Parameterization of the planetary boundary layer in the UCLA general circulation model: Formulation and results. *Mon. Wea. Rev.*, **111**, 2224-2243.